

1. Let M be a finitely generated R -module, set $J = J(R)$, the Jacobson radical of R . Prove: if $M = MJ$ then $M = 0$. Is this true if M is not finitely generated?
2. (i) Prove that an integral extension of a Jacobson ring is Jacobson.
(ii) Prove that the polynomial ring $F[t]$ over a field F is a Jacobson ring
(iii) Find a PID R such that $J(R) \neq 0$. Can you complete the sentence: A PID is a Jacobson ring if and only if ...
3. Complete the missing details in the proof of Proposition 41: show that the maps c and e respect inclusions and intersections of ideals, and that e respects sums.
4. (i) Let $f(t_1, \dots, t_n)$ be a polynomial over a field F . Suppose there exist infinite subsets X_1, \dots, X_n of F such that $f(x_1, \dots, x_n) = 0$ for all $(x_1, \dots, x_n) \in X_1 \times \dots \times X_n$. Prove that f is the zero polynomial.
(ii) Let U be an algebraic set in F^n and V an algebraic set in F^m . Show that $U \times V$ is an algebraic set in F^{n+m} . A subset X of U is dense in U if U is the smallest algebraic set that contains X . Show that if X is a dense subset of U and Y is a dense subset of V then $X \times Y$ is dense in $U \times V$. How does this relate to (i)?
5. (i) Let M be a finitely generated R -module and $\phi : M \rightarrow M$ a module endomorphism. Prove that if ϕ is surjective then ϕ is an isomorphism). (Hint: Consider M as an $R[t]$ -module where t acts like ϕ .)
(ii) Let $F = R^d$ be a free module and Y a generating set for F . Prove that if $|Y| \leq d$ then Y is a basis and $|Y| = d$.
6. A module M has length $\lambda(M) = n$ if there is a chain of submodules $0 = M_0 < M_1 < \dots < M_n = M$ and n is maximal, $\lambda(M) = \infty$ if there is no maximal such n .
(i) Prove that length is additive on extensions of modules, i.e. if $N \leq M$ then $\lambda(M) = \lambda(N) + \lambda(M/N)$.
(ii) Suppose that M is a Noetherian R -module and that $MP^k = 0$ for some maximal ideal P of R and some integer k . Show that M has finite length.