

Gravitational Collapse for Newtonian stars

M.R.I. SCHRECKER

*Department of Mathematics
University College London*

Classical model of a star: sphere of gas under Newtonian gravity.

- Balance between pressure and gravity in a static star;
- As gas burns, balance shifts;

Classical model of a star: sphere of gas under Newtonian gravity.

- Balance between pressure and gravity in a static star;
- As gas burns, balance shifts;
- Possible collapse? Supernova?




Figure: Image credit: R.J. Hall

The Euler-Poisson equations of gas dynamics with Newtonian gravity:

$$\begin{cases} \partial_t \rho + \operatorname{div}_{\mathbf{x}}(\rho \mathbf{u}) = 0, & (t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^3, \\ \rho(\partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u}) + \nabla_{\mathbf{x}} p(\rho) = -\rho \nabla \Phi, & (t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^3, \\ \Delta \Phi = 4\pi \rho, & (t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^3. \end{cases} \quad (1)$$

ρ is density, \mathbf{u} is velocity, p is pressure, Φ is gravitational potential.

We assume the equation of state

$$p = p(\rho) = \rho^\gamma, \quad \gamma \in (1, \frac{4}{3}).$$

The Euler-Poisson equations of gas dynamics with Newtonian gravity:

$$\begin{cases} \partial_t \rho + \operatorname{div}_{\mathbf{x}}(\rho \mathbf{u}) = 0, & (t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^3, \\ \rho(\partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u}) + \nabla_{\mathbf{x}} p(\rho) = -\rho \nabla \Phi, & (t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^3, \\ \Delta \Phi = 4\pi \rho, & (t, \mathbf{x}) \in \mathbb{R} \times \mathbb{R}^3. \end{cases} \quad (1)$$

ρ is density, \mathbf{u} is velocity, p is pressure, Φ is gravitational potential.
We assume the equation of state

$$p = p(\rho) = \rho^\gamma, \quad \gamma \in (1, \frac{4}{3}).$$



Example adiabatic exponents

$\gamma = \frac{5}{3}$ – monatomic gas, used for fully convective star cores (e.g. red giants);
 $\gamma = \frac{4}{3}$ – high mass white dwarf stars, main-sequence stars (e.g. the Sun).
In general, as γ decreases, density is increasingly weighted towards centre.

Collapse is the formation of a *singularity* at the origin, i.e.

$$\rho(t, 0) \rightarrow \infty \quad \text{as} \quad t \rightarrow 0 - .$$

- For $\gamma > \frac{4}{3}$, no finite mass and energy collapse possible.
- For $\gamma = \frac{4}{3}$, Goldreich–Weber collapse - unsuitable model for outer core.




Figure: GIF credit: NASA

Self-similarity and singularities interact in a wide range of problems.

- Stellar collapse;
- Formation/expansion of shock waves;
- Shock reflection;
- Droplet pinch-off;
- Bacterial growth;
- Geometric wave equations;
- Yang–Mills;
- ...

Self-similarity and singularities interact in a wide range of problems.

- Stellar collapse;
- Formation/expansion of shock waves;
- Shock reflection;
- Droplet pinch-off;
- Bacterial growth;
- Geometric wave equations;
- Yang–Mills;
- ...

Key Features:

- Non-linearity;
- Intertwining of spatial and time scales;
- Good initial data leads to badly behaved solutions!

Scaling

Let $\rho = \rho(t, r)$, $\mathbf{u} = u(t, r) \frac{\mathbf{x}}{|\mathbf{x}|}$, $r = |\mathbf{x}|$, solve Euler-Poisson, $\lambda > 0$.
Then

$$\rho_\lambda(t, r) = \lambda^{-\frac{2}{2-\gamma}} \rho\left(\frac{t}{\lambda^{\frac{1}{2-\gamma}}}, \frac{r}{\lambda}\right), \quad u_\lambda(t, r) = \lambda^{-\frac{\gamma-1}{2-\gamma}} u\left(\frac{t}{\lambda^{\frac{1}{2-\gamma}}}, \frac{r}{\lambda}\right)$$

is also a solution. (NB: This is a *unique* scaling!)

Scaling

Let $\rho = \rho(t, r)$, $\mathbf{u} = u(t, r) \frac{\mathbf{x}}{|\mathbf{x}|}$, $r = |\mathbf{x}|$, solve Euler-Poisson, $\lambda > 0$.
Then

$$\rho_\lambda(t, r) = \lambda^{-\frac{2}{2-\gamma}} \rho\left(\frac{t}{\lambda^{\frac{1}{2-\gamma}}}, \frac{r}{\lambda}\right), \quad u_\lambda(t, r) = \lambda^{-\frac{\gamma-1}{2-\gamma}} u\left(\frac{t}{\lambda^{\frac{1}{2-\gamma}}}, \frac{r}{\lambda}\right)$$

is also a solution. (NB: This is a *unique* scaling!)

Self-similarity

We define a *self-similar* variable

$$y = \frac{r}{(-t)^{2-\gamma}},$$

and search for

$$\rho(t, r) = (-t)^{-2} \tilde{\rho}(y), \quad u(t, r) = (-t)^{1-\gamma} \tilde{u}(y).$$

Natural notions of mass and energy for Euler-Poisson:

$$M[\rho] = \int_0^\infty \rho r^2 dr, \quad E[\rho, u] = \int_0^\infty \left(\rho u^2 + \frac{\rho^\gamma}{\gamma - 1} + \frac{1}{2} \rho \Phi \right) r^2 dr,$$

where Φ solves $\Delta \Phi = 4\pi \rho$ is the gravitational potential.

Under scaling,

$$M[\rho_\lambda] = \lambda^{\frac{4-3\gamma}{2-\gamma}} M[\rho], \quad E[\rho_\lambda, u_\lambda] = \lambda^{\frac{6-5\gamma}{2-\gamma}} E[\rho, u].$$

Thus $\gamma = \frac{4}{3}$ is *mass-critical*, $\gamma = \frac{6}{5}$ is *energy-critical*.

Defining a convenient variable $\omega(y) = \tilde{u}(y)/y + 2 - \gamma$, self-similar Euler-Poisson becomes

$$\begin{aligned}\tilde{\rho}' &= \frac{y\tilde{\rho}h(\tilde{\rho}, \omega)}{\gamma\tilde{\rho}^{\gamma-1} - y^2\omega^2}, \\ \omega' &= \frac{4 - 3\gamma - 3\omega}{y} - \frac{y\omega h(\tilde{\rho}, \omega)}{\gamma\tilde{\rho}^{\gamma-1} - y^2\omega^2},\end{aligned}\tag{2}$$

where $h(\tilde{\rho}, \omega)$ is a quadratic function.

Defining a convenient variable $\omega(y) = \tilde{u}(y)/y + 2 - \gamma$, self-similar Euler-Poisson becomes

$$\begin{aligned}\tilde{\rho}' &= \frac{y\tilde{\rho}h(\tilde{\rho}, \omega)}{\gamma\tilde{\rho}^{\gamma-1} - y^2\omega^2}, \\ \omega' &= \frac{4 - 3\gamma - 3\omega}{y} - \frac{y\omega h(\tilde{\rho}, \omega)}{\gamma\tilde{\rho}^{\gamma-1} - y^2\omega^2},\end{aligned}\tag{2}$$

where $h(\tilde{\rho}, \omega)$ is a quadratic function.

Definition (Sonic point)

Let $(\tilde{\rho}(\cdot), \omega(\cdot))$ be a C^1 -solution to the self-similar Euler-Poisson system on the interval $(0, \infty)$. A point $y_* \in (0, \infty)$ such that

$$G(y, \tilde{\rho}, \omega) := \gamma\tilde{\rho}^{\gamma-1}(y_*) - y_*^2\omega^2(y_*) = 0$$

is called a *sonic point*.

Initial/boundary conditions

For a regular solution, we require

$$\tilde{\rho}(0) > 0, \quad \omega(0) = \frac{4 - 3\gamma}{3},$$
$$\tilde{\rho}(y) \sim y^{-\frac{2}{2-\gamma}} \text{ as } y \rightarrow \infty, \quad \lim_{y \rightarrow \infty} \omega(y) = 2 - \gamma.$$

NB: this forces the existence of a sonic point!

Initial/boundary conditions

For a regular solution, we require

$$\tilde{\rho}(0) > 0, \quad \omega(0) = \frac{4 - 3\gamma}{3},$$

$$\tilde{\rho}(y) \sim y^{-\frac{2}{2-\gamma}} \text{ as } y \rightarrow \infty, \quad \lim_{y \rightarrow \infty} \omega(y) = 2 - \gamma.$$

NB: this forces the existence of a sonic point!

Theorem (Guo–Hadzic–Jang–S. '21)

For each $\gamma \in (1, \frac{4}{3})$, there exists a global, real-analytic solution $(\tilde{\rho}, \omega)$ of self-similar Euler-Poisson with a single sonic point y_ such that:*

$$\tilde{\rho}(y) > 0 \text{ for all } y \in [0, \infty), \quad -\frac{2}{3}y < u(y) < 0 \text{ for all } y \in (0, \infty).$$

In addition, both ρ and ω are strictly monotone:

$$\tilde{\rho}'(y) < 0 \text{ for all } y \in (0, \infty), \quad \omega'(y) > 0 \text{ for all } y \in (0, \infty).$$

Classical and numerical work

- Taylor, Von Neumann, Sedov, Gülderley '40s: study implosion and explosion for Euler equations;
- Larson–Penston '69: numerical solution for $\gamma = 1$;
- Hunter '77: family of numerical solutions for $\gamma = 1$;
- Yahil '83: numerical solutions for $\gamma \in [\frac{6}{5}, \frac{4}{3})$;
- Maeda–Harada '01: numerical evidence towards mode stability of Larson–Penston.

Classical and numerical work

- Taylor, Von Neumann, Sedov, Gülderley '40s: study implosion and explosion for Euler equations;
- Larson–Penston '69: numerical solution for $\gamma = 1$;
- Hunter '77: family of numerical solutions for $\gamma = 1$;
- Yahil '83: numerical solutions for $\gamma \in [\frac{6}{5}, \frac{4}{3})$;
- Maeda–Harada '01: numerical evidence towards mode stability of Larson–Penston.

Recent works

- Merle–Raphaël–Rodnianski–Szeftel '19: existence of a imploding self-similar solutions for Euler;
- Guo–Hadzic–Jang '20: construction of LP solution.

Regularity

Expect stability tied to regularity (MRRS '19). Requires smoothness through sonic point.

Regularity

Expect stability tied to regularity (MRRS '19). Requires smoothness through sonic point.

Non-linearity

Methods need to be adapted to specific non-linearities (no general recipe for solving such problems).

Regularity

Expect stability tied to regularity (MRRS '19). Requires smoothness through sonic point.

Non-linearity

Methods need to be adapted to specific non-linearities (no general recipe for solving such problems).

Non-autonomous system

Non-autonomous forces evolving phase portrait. No fixed phase portrait analysis for invariant regions.

Two explicit solutions

Far-field solution (ρ_f, ω_f) and Friedman solution (ρ_F, ω_F) :

$$(\rho_f(y), \omega_f(y)) = (k_\gamma y^{-\frac{2}{2-\gamma}}, 2 - \gamma), \quad (\rho_F(y), \omega_F(y)) = \left(\frac{1}{6\pi}, \frac{4}{3} - \gamma\right).$$

Two explicit solutions

Far-field solution (ρ_f, ω_f) and Friedman solution (ρ_F, ω_F) :

$$(\rho_f(y), \omega_f(y)) = (k_\gamma y^{-\frac{2}{2-\gamma}}, 2 - \gamma), \quad (\rho_F(y), \omega_F(y)) = \left(\frac{1}{6\pi}, \frac{4}{3} - \gamma\right).$$

- Sonic points at $y_f(\gamma) < y_F(\gamma)$.
- Far-field satisfies asymptotic boundary condition as $y \rightarrow \infty$.
- Friedman satisfies boundary condition at origin.

Idea: Use $\omega_f = 2 - \gamma$, $\omega_F = \frac{4}{3} - \gamma$ as barriers.

Proposition (Local Solution)

For all $\gamma \in (1, \frac{4}{3})$, there exists $\nu > 0$ such that for all $y_ \in [y_f(\gamma), y_F(\gamma)]$, there exists an analytic solution $(\rho(\cdot; y_*), \omega(\cdot; y_*))$ to self-similar Euler-Poisson on $(y_* - \nu, y_* + \nu)$ with a single sonic point at y_* .*

Idea: By formal Taylor expansion and selection of physical first order coefficient


Proposition (Local Solution)

For all $\gamma \in (1, \frac{4}{3})$, there exists $\nu > 0$ such that for all $y_ \in [y_f(\gamma), y_F(\gamma)]$, there exists an analytic solution $(\rho(\cdot; y_*), \omega(\cdot; y_*))$ to self-similar Euler-Poisson on $(y_* - \nu, y_* + \nu)$ with a single sonic point at y_* .*

Idea: By formal Taylor expansion and selection of physical first order coefficient

Lemma (Solving to the right)

For each $\gamma \in (1, \frac{4}{3})$, each $y_ \in [y_f(\gamma), y_F(\gamma)]$, the local solution $(\rho(\cdot; y_*), \omega(\cdot; y_*))$ obtained by Taylor expansion extends globally to the right on $[y_*, \infty)$, remains supersonic, and satisfies the asymptotic boundary conditions.*



Key ideas


- Use structure of $h(\rho, \omega)$ and $G(y; \rho, \omega)$ to derive dynamical invariances to the right.
- Show ω remains trapped between $\frac{4}{3} - \gamma$ and $2 - \gamma$.
- Extend dynamical invariance to show flow remains supersonic.
- Asymptotics follow easily from structure of flow.

Aim: Find \bar{y}_* such that local solution $(\rho(\cdot; \bar{y}_*), \omega(\cdot; \bar{y}_*))$ extends smoothly to $y = 0$. Look for solution with

$$\frac{4}{3} - \gamma \leq \omega(y; \bar{y}_*) < 2 - \gamma, \quad \lim_{y \rightarrow 0} \omega(y; \bar{y}_*) = \frac{4}{3} - \gamma.$$


Aim: Find \bar{y}_* such that local solution $(\rho(\cdot; \bar{y}_*), \omega(\cdot; \bar{y}_*))$ extends smoothly to $y = 0$. Look for solution with

$$\frac{4}{3} - \gamma \leq \omega(y; \bar{y}_*) < 2 - \gamma, \quad \lim_{y \rightarrow 0} \omega(y; \bar{y}_*) = \frac{4}{3} - \gamma.$$



Aim: Find \bar{y}_* such that local solution $(\rho(\cdot; \bar{y}_*), \omega(\cdot; \bar{y}_*))$ extends smoothly to $y = 0$. Look for solution with

$$\frac{4}{3} - \gamma \leq \omega(y; \bar{y}_*) < 2 - \gamma, \quad \lim_{y \rightarrow 0} \omega(y; \bar{y}_*) = \frac{4}{3} - \gamma.$$



Try: $\bar{y}_* = \inf Y = \inf \left\{ y_* \in (y_f, y_F) \mid \exists y \text{ such that } \omega(y; y_*) = \frac{4 - 3\gamma}{3} \right\}.$

Key idea: Prove monotonicity for both $\rho(\cdot; y_*)$ and $\omega(\cdot; y_*)$ as long as $y_* \in Y$ and $\omega(\cdot; y_*) \geq \frac{4}{3} - \gamma$.

Linear Stability

- Appropriate self-similar coordinates;
- Non-self-adjoint problem (complex eigenvalues);
- Sonic degeneracy and issues with dissipativity (monotonicity).

Linear Stability

- Appropriate self-similar coordinates;
- Non-self-adjoint problem (complex eigenvalues);
- Sonic degeneracy and issues with dissipativity (monotonicity).

Future directions

- Non-linear stability;
- Einstein-Euler (relativistic self-similar fluid implosion) and its stability (cf. Guo–Hadžić–Jang '21).
- Continuation and expansion?

Thank you!

