

## Warm-up

1. Differentiate  $x^{17} - x^{-17}$  with respect to  $x$ .
  - The derivative of  $x^a$  is  $ax^{a-1}$  so the derivative of this expression is  $17x^{16} + 17x^{-18}$ .
2. Differentiate  $2\sqrt{x} + 3\sqrt[3]{x}$  with respect to  $x$ .
  - Remember that  $\sqrt{x} = x^{1/2}$  and  $\sqrt[3]{x} = x^{1/3}$ , so the derivative of this expression is  $x^{-1/2} + x^{-2/3}$ , which we could write as  $\frac{1}{\sqrt{x}} + \frac{1}{x^{2/3}}$  if we wanted to.
3. Differentiate  $1 - e^{3x}$  with respect to  $x$ .
  - Remember that the derivative of a constant is 0, so the derivative of this expression is just  $-3e^{3x}$ .
4. Find the tangent to the curve  $y = e^x + x^2$  at  $x = 2$ .
  - We need to find the value of the derivative  $\frac{dy}{dx}$  at  $x = 2$  because that's equal to the gradient of the tangent. We can differentiate to find  $\frac{dy}{dx} = e^x + 2x$  so that gradient we wanted is  $e^2 + 4$ . We also want the tangent to have the same value at  $x = 2$  as the curve; that's  $e^x + x^2$  at  $x = 2$ , which is also  $e^2 + 4$ . So our tangent is  $y = (e^2 + 4)(x - 1)$ .
5. Find the normal to the parabola  $y = x^2$  at  $x = 3$ .
  - First find the derivative at  $x = 3$ , which is 6 for this parabola. That's the gradient of the tangent, and the normal is at right-angles to the tangent, so it has gradient  $-\frac{1}{6}$ . We have  $y = -\frac{x}{6} + c$  and we want the normal to go through the point  $(3, 9)$ . So we want  $c = \frac{19}{2}$ .
6. Find the turning points of the curve  $y = x^4 - 2x^3 + x^2$ . Identify whether the turning points are maxima or minima. For which values of  $x$  is  $y = x^4 - 2x^3 + x^2$  increasing? For which values of  $x$  is it decreasing?
  - The turning points must have  $\frac{dy}{dx} = 0$  so we must have  $4x^3 - 6x^2 + 2x = 0$ . That happens when  $x = 0$  or when  $2x^2 - 3x + 1 = 0$  which happens when  $(2x - 1)(x - 1) = 0$ , which is either  $x = 1$  or  $x = \frac{1}{2}$ .
  - Now find the second derivative to check whether these are minima or maxima.  $\frac{d^2y}{dx^2} = 12x^2 - 12x + 2$ , which is positive for  $x = 0$ , negative for  $x = \frac{1}{2}$ , and positive for  $x = 1$ . So we have a (local) minimum, then a (local) maximum, then a (local) minimum.
  - The function is decreasing for  $x < 0$ , then increasing for  $0 < x < \frac{1}{2}$ , then decreasing for  $\frac{1}{2} < x < 1$  then increasing for  $x > 1$ .

7. Two points  $A$  and  $B$  are on the curve  $y = x^3 + x^2 + x + 1$ .  $A$  is held fixed at  $(1, 4)$ . The point  $B$  is moved along the curve towards  $A$ . What happens to the line through  $A$  and  $B$ ?

- The line definitely goes through  $A$ , which doesn't move. The thing we learn from "differentiation from first principles" is that the gradient of the line gets closer and closer to the derivative of the function at  $A$ .
- The derivative is  $3x^2 + 2x + 1$  which is 6 at  $x = 1$ . The value is 4, so the tangent is  $y = 6x - 2$ . So if the line through  $A$  and  $B$  is  $y = mx + c$  then  $m$  gets closer and closer to 6 and  $c$  gets closer and closer to  $-2$ .

8. Suppose that the derivative of a polynomial  $p(x)$  with respect to  $x$  is  $q(x)$ . Find an expression for  $\int q(x) dx$ .

- This must be  $p(x) + c$  for some constant  $c$ .

9. Find the area enclosed between the polynomial  $y = x^2 + 4x + 3$  and the  $x$ -axis.

- First find the points where  $y = 0$ . We have  $(x + 3)(x + 1) = 0$  so these points are at  $x = -1$  or  $x = -3$ . In between, we have  $y < 0$  (by considering the graph).
- So we want  $-\int_{-3}^{-1} x^2 + 4x + 3 dx$ . That minus sign out the front is because the function is negative.
- This works out to be  $\frac{4}{3}$ .

10. Find  $\int_{-1}^1 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 dx$ .

- This is  $\left[ x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \frac{x^5}{5} + \frac{x^6}{6} + \frac{x^7}{7} \right]_{-1}^1 = 2 + \frac{2}{3} + \frac{2}{5} + \frac{2}{7} = \frac{352}{105}$ .

11. Find  $\int_{-\pi}^{\pi} x^{1729} dx$ .

- This is  $\left[ \frac{x^{1730}}{1730} \right]_{-\pi}^{\pi} = 0$ . In general, any "odd function" with  $f(-x) = -f(x)$  which we integrate from  $-a$  to  $a$  for any real number  $a$  will give zero.

12. Find

$$\int \frac{x+3}{x^3} dx, \quad \int \sqrt[3]{x} dx, \quad \int ((x^2)^3)^5 dx, \quad \int (x^2+1)^3 dx$$

- $\int \frac{x+3}{x^3} dx = \int \frac{1}{x^2} + \frac{3}{x^3} dx = -\frac{1}{x} - \frac{3}{2x^2} + c$  where  $c$  is a constant.

- $\int \sqrt[3]{x} \, dx = \int x^{1/3} \, dx = \frac{3}{4}x^{4/3} + c$  where  $c$  is a constant.
- $\int \left((x^2)^3\right)^5 \, dx = \int x^{30} \, dx = \frac{x^{31}}{31} + c$  where  $c$  is a constant.
- $\int (x^2 + 1)^3 \, dx = \int x^6 + 3x^4 + 3x^2 + 1 \, dx = \frac{x^7}{7} + \frac{3x^5}{5} + x^3 + x + c$  where  $c$  is a constant.

13. By thinking about the area that the integral represents, explain why

$$\int_{-1}^1 f(x) \, dx = \int_{-1}^1 f(-x) \, dx.$$

- The graph of  $f(-x)$  is the graph of  $f(x)$  reflected in  $y$ -axis. Also, note that if we reflect the interval  $-1 \leq x \leq 1$  in the  $y$ -axis then we get the same interval back. On the left-hand side, we're finding the area under  $f(x)$  (or maybe negative the area in any regions where  $f$  is negative). On the right-hand side, we're calculating exactly the same area, but with the shape of the graph reflected.

## MAT questions

### MAT 2014 Q1C

- First work out the derivative  $3kx^2 - 2(k+1)x + (2-k)$  and then the second derivative  $6kx - 2(k+1)$ .
- At  $x = 1$ , the first derivative is  $3k - 2(k+1) + (2-k)$ , which is zero for any  $k$ , and the second derivative is  $6k - 2(k+1) = 4k - 2$ .
- For this to be a local minimum we would need zero derivative (which we have!) and we would need the second derivative to be positive, so we would need  $k > \frac{1}{2}$ .
- The answer is (c).

**MAT 2017 Q1A**

- Let's look for the stationary points by working out the derivative  $6x^2 - 2kx + 2$ .
- That's got two distinct roots if the discriminant  $4k^2 - 48$  is positive, which happens if  $k^2 > 12$ .
- Technically we should probably check that the second derivative isn't zero at these points, but since the first derivative is a quadratic, it can only have zero derivative at a root if there's a single repeated root (which is not the case).
- The answer is (b).

**MAT 2014 Q1J**

- This question is tough. We're not going to work out what the function  $f(x)$  is! We're just going to work out precisely the information  $\int_{-1}^1 f(x) dx$ . Let's give that number a name (it is a number, not a function of  $x$ ). I'm going to call it  $A$ .
- Now the equation we've been given is

$$6 + f(x) = 2f(-x) + 3Ax^2$$

- I don't know much about  $f(x)$ , and frankly I don't care much about the details of  $f(x)$ , I just want to find  $A$ . I've noticed that if I integrate the left-hand side of the equation, I get  $12 + A$ .
- If I integrate the right-hand side I get


$$2 \int_{-1}^1 f(-x) dx + 3A \int_{-1}^1 x^2 dx.$$

I've moved the  $A$  outside of the integral there because it's just a constant.

- Now note that  $\int_{-1}^1 f(-x) dx = \int_{-1}^1 f(x) dx$ ; we talked about this in the warm-up. That's  $A$ .
- So we have  $12 + A = 2A + 3A \int_{-1}^1 x^2 dx$ . We're almost done, I just need to calculate one integral and I'll get an equation for  $A$ .
- I get  $12 + A = 2A + 2A$  so  $A = 4$ .
- The answer is (a).

**MAT 2016 Q1H**

- First, let's calculate the two areas in the question.



- The  $x$ -axis and  $f(x)$  enclose a region with  $-\sqrt{a} < x < \sqrt{a}$  where  $f(x)$  is positive. The area is

$$\int_{-\sqrt{a}}^{\sqrt{a}} a - x^2 \, dx = 2a^{3/2} - \frac{2}{3}a^{3/2} = \frac{4}{3}a^{3/2}.$$

- The  $x$ -axis and  $g(x)$  enclose a region with  $-a^{1/4} < x < a^{1/4}$  where  $g(x)$  is negative. The area is

$$-\int_{-a^{1/4}}^{a^{1/4}} x^4 - a \, dx = -\frac{2}{5}a^{5/4} + 2a^{5/4} = \frac{8}{5}a^{5/4}$$


- We want to find out when the first area is bigger. We need

$$\frac{4}{3}a^{3/2} > \frac{8}{5}a^{5/4}.$$

- This rearranges to  $a > \left(\frac{6}{5}\right)^4$ .
- The answer is (e).

**MAT 2017 Q3**

(i) Here's a sketch



Note that all the curves pass through the point  $(0,0)$  and through  $(1,1)$ . For  $0 < x < 1$  the curve with the highest value of  $k$  is the largest, but for  $x > 1$  the curve with the lowest value of  $k$  is the largest.


(ii) For  $0 < x < 1$ , the curve  $y = f_{k+1}(x)$  has a larger value than the curve  $y = f_k(x)$ . The area we want is

$$\int_0^1 f_{k+1}(x) - f_k(x) \, dx = \int_0^1 x^{1/(k+1)} - x^{1/k} \, dx = \left[ \frac{x^{1+1/(k+1)}}{1+1/(k+1)} - \frac{x^{1+1/k}}{1+1/k} \right]_0^1 = \frac{k+1}{k+2} - \frac{k}{k+1}.$$

We can check that if  $k = 1$  then this is  $\frac{2}{3} - \frac{1}{2} = \frac{1}{6}$ .

(iii) Remember that  $y = f_1(x)$  is just  $y = x$ . This intersects the line  $y = c$  for  $x = c$ , so the point of intersection is  $(c, c)$ . The curve  $y = f_2(x)$  is just  $y = x^{1/2}$  so the point of intersection with the line  $y = c$  is  $(c^2, c)$ .

(iv) Here's a sketch



We can write the area between the curves and under  $y = c$  in two parts as

$$A = \int_0^{c^2} x^{1/2} - x \, dx + \int_{c^2}^c c - x \, dx.$$

We can calculate these integrals to get (after simplifying)

$$A = \frac{c^2}{2} - \frac{c^3}{3}.$$

Now for the two regions to have equal area, they must each have area  $\frac{1}{12}$ , because we found in part (ii) that the overall area is  $\frac{1}{6}$ . This gives the equation  $\frac{1}{12} = \frac{c^2}{2} - \frac{c^3}{3}$  which we can rearrange for  $4c^3 - 6c^2 + 1 = 0$ .

We know that  $0 < c < 1$ , so let's try guessing simple fractions as roots of this cubic. It turns out that  $c = \frac{1}{2}$  works. Perhaps we could factorise the cubic to check that this is the only root in that range, but it's easier to argue geometrically; as  $c$  increases from zero to one, the area under the line  $y = c$  only increases, so we can't have two different values of  $c$  which split the region in half.

## Reflection

- Splitting the area up into multiple parts is a good tactic. In the long question, I split the area up into a region to the left of a point of intersection and a separate region to the right. I could have spotted that the region to the right was just a right-angled isosceles triangle, which might have been a faster way to get the area.
- If you've never seen something like  $\int_{-1}^1 f(t) \, dt$  before, it's worth thinking about what this means and why it's the same thing as  $\int_{-1}^1 f(x) \, dx$ .
- Drawing sketches is still important, even if the question doesn't ask you to!