

# On global solutions to the Navier-Stokes equations

Isabelle Gallagher

Institut de Mathématiques de Jussieu, Université Paris-Diderot

Oxford, September 2012

# Presentation of the equations

- Viscous, incompressible, homogeneous fluid, in  $\mathbb{R}^3$
- Velocity  $u = (u^1, u^2, u^3)(t, x)$ , pressure  $p(t, x)$

$$(NS) \quad \begin{cases} \partial_t u + u \cdot \nabla u - \Delta u = -\nabla p \\ \operatorname{div} u = 0 \end{cases}$$

with

$$\Delta u = \sum_{j=1}^3 \partial_j^2 u, \quad \operatorname{div} u = \sum_{j=1}^3 \partial_j u^j, \quad \partial_j := \frac{\partial}{\partial x_j}, \quad \partial_t := \frac{\partial}{\partial t}$$
$$u \cdot \nabla u = \sum_{j=1}^3 u^j \partial_j u = \sum_{j=1}^3 \partial_j (u^j u).$$

**Remark :** The pressure can be eliminated by **projection onto divergence-free vector fields** :  $\mathbb{P} = \operatorname{Id} - \nabla \Delta^{-1} \operatorname{div}$ .

**Cauchy data :**  $u|_{t=0} = u_0$ .

# Solving the equations

We want to find  $u(t, x)$  solution to (NS) in some sense (distributional, classical...), such that  $u(0, x) \equiv u_0(x)$ .

Standard methods :

- **Compactness** methods :

- Find an *a priori* bound on the solution :  $\|u(t)\|_X \leq C(u_0)$  ;
- Construct a sequence of *approximate equations*  $(NS)_{n \in \mathbb{N}}$  which can be solved by the Cauchy-Lipschitz theorem : this yields a sequence of *approximate solutions*  $(u_n)_{n \in \mathbb{N}}$ , uniformly bounded in  $X$  ;
- Use the uniform bound in  $X$  to construct weak limit points to the sequence  $(u_n)_{n \in \mathbb{N}} : u_n \rightharpoonup u$  ;
- Use space-time compactness to prove that  $u$  solves (NS).

# Solving the equations

- Banach **fixed point** theorem :
  - Write the equation in integral form :

$$u(t) = e^{t\Delta} u(0) + B(u, u)(t)$$

- Apply a fixed point theorem.

# Fundamental properties of (NS) (I)

- **Conservation of the energy**

Conservation of energy is due to the formal identity

$$\frac{1}{2} \|u(t)\|_{L^2}^2 + \int_0^t \|\nabla u(t')\|_{L^2}^2 dt' = \frac{1}{2} \|u_0\|_{L^2}^2$$

thanks to the structure of the nonlinear term :

$$(\mathbb{P}(u \cdot \nabla u) | u)_{L^2} = 0.$$

So in particular  $u \in L^\infty(\mathbb{R}^+; L^2)$  and  $\nabla u \in L^2(\mathbb{R}^+; L^2)$ .

# Fundamental properties of (NS) (II)

- **Scale invariance**

If  $u(t, x)$  is a solution of (NS) associated with the initial data  $u_0(x)$  on  $[0, T] \times \mathbb{R}^3$ , then for all  $\lambda > 0$ ,  $a \in \mathbb{R}^3$

$$u_\lambda(t, x) := \lambda u(\lambda^2 t, \lambda(x - a))$$

is a solution associated with  $u_{\lambda,0}(x) := \lambda u_0(\lambda(x - a))$  on  $[0, \lambda^{-2} T] \times \mathbb{R}^3$ .

# Weak solutions

Using the **conservation of energy**, one can prove the following result.

**Theorem** [Leray, 1934]

Let  $u_0 \in L^2(\mathbb{R}^d)$  be a divergence free vector field. There is a solution  $u$  of (NS) satisfying for all  $t \geq 0$

$$\|u(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla u(t')\|_{L^2}^2 dt' \leq \|u_0\|_{L^2}^2.$$

Remarks :

- ▶ Proof by **compactness**.
- ▶ Search for conditions on the initial data to guarantee uniqueness (if  $d = 2$ , OK — due to scale invariance).

# Strong solutions

One does not use the **structure** of the equation, but rather its **scale invariance**, by a **fixed point** method.

Solving (NS) is equivalent to solving

$$u = e^{t\Delta} u_0 + \mathbb{B}(u, u)$$

where  $e^{t\Delta}$  is the heat semi-group on  $\mathbb{R}^d$  and  $\mathbb{B}$  the bilinear form

$$\mathbb{B}(u, u)(t) := - \int_0^t e^{(t-t')\Delta} \mathbb{P} \operatorname{div} (u \otimes u)(t') dt' .$$

The problem consists in finding an **adapted** Banach space  $X$ , such that  $\mathbb{B}$  is continuous from  $X \times X$  to  $X$ .



# An existence and uniqueness result

## Theorem

Let  $X$  be an adapted space. If  $u_0$  is such that  $\|e^{t\Delta} u_0\|_X$  is small enough, then there is a unique solution to (NS) in  $X$ .

**Remarks :** • By scale invariance, the norm on  $X$  must satisfy

$$\forall \lambda > 0, \forall x \in \mathbb{R}^3, \quad \lambda \|f(\lambda^2 t, \lambda(x - a))\|_X \sim \|f\|_X.$$

- This corresponds to **small initial data** or **small time** results.

## Examples :

- Leray '34 : smallness measured by  $\|u_0\|_{L^2} \|\nabla u_0\|_{L^2}$  if  $d = 3$
- Fujita-Kato '64 with  $\|u_0\|_{\dot{H}^{\frac{d}{2}-1}}$
- Kato '84 with  $\|u_0\|_{L^d}$
- Cannone-Meyer-Planchon '94 with  $\|u_0\|_{B_p}$ , where

$$\|u_0\|_{B_p} := \sup_{t>0} t^{\frac{1}{2}(1-\frac{d}{p})} \|e^{t\Delta} u_0\|_{L^p}.$$

# The optimal adapted space

Recall

$$\|u_0\|_{B_p} := \sup_{t>0} t^{\frac{1}{2}(1-\frac{d}{p})} \|e^{t\Delta} u_0\|_{L^p}.$$

- Any Banach space of tempered distributions, scale and translation invariant, is **embedded in**  $B_\infty$  [Meyer '96]
- (NS) is **ill-posed** in  $B_\infty$  [Bourgain-Pavlovic '08, Germain '08]
- (NS) is **well-posed** (for small enough data) in  $\tilde{B}_\infty$  where

$$\|u_0\|_{\tilde{B}_\infty} := \|u_0\|_{B_\infty} + \sup_{\substack{x \in \mathbb{R}^d \\ R > 0}} \frac{1}{R^{\frac{d}{2}}} \left( \int_{P(x,R)} |(e^{t\Delta} u_0)(t,y)|^2 dy dt \right)^{\frac{1}{2}}$$

with  $P(x, R) := [0, R^2] \times B(x, R)$  [Koch-Tataru '01].

## Remarks (I)

In this context in general, only **small data** or **small time** theorems are known. They hold (with the same proof) for the more general equation

$$\partial_t u - \Delta u = Q(u, u)$$

where  $Q(v, w) := \sum_{1 \leq j, k \leq 3} Q_{j,k}(D)(v^j w^k)$  and  $Q_{j,k}(D)$  are smooth homogeneous Fourier multipliers of order 1.

However some of these equations are known to produce **blow-up in finite time** [Montgomery-Smith '01], including for (large) data for which Navier-Stokes does not [G-Paicu '09].

## Remarks (II)

There is a **discrepancy** between the energy (providing control of norms) and the scaling (necessary to implement the fixed point).

If  $d = 2$ , the energy space is scale invariant, the equation is said **critical**.

In dimension  $d \geq 3$ , there are  $d/2 - 1$  derivatives between scaling and energy : the equation is said **supercritical**.

# Properties of $\mathcal{G}$

In the following we denote by  $\mathcal{G}$  the space of initial data generating a **global smooth solution** to the three-dimensional Navier-Stokes equations.

We want to study **geometrical** properties of  $\mathcal{G}$ .

We shall prove that  $\mathcal{G}$  is

- **open (strong topology)** in  $B_p$  [G-Iftimie-Planchon '03],  $BMO^{-1}$  [Auscher, Dubois, Tchamitchian '04]
- **connected** in  $\dot{H}^{\frac{1}{2}}$ ,  $B_p$  [G-Iftimie-Planchon '03],  $BMO^{-1}$  [Auscher, Dubois, Tchamitchian '04]
- **unbounded** in  $B_\infty$  [Chemin-G '06,'09,'10, Chemin-G-Paicu '12, Chemin-G-Zhang '12]
- **open (weak topology)** (under an anisotropy assumption) [Bahouri-G '12, Bahouri-Chemin-G in progress].

# The set $\mathcal{G}$ is **strongly** open, and connected

Let us prove the following result.

**Theorem** [G-Iftimie-Planchon '03]

Let  $u \in C(\mathbb{R}^+, \dot{H}^{\frac{1}{2}}(\mathbb{R}^3))$  be a solution to (NS). Then

$$\lim_{t \rightarrow +\infty} \|u(t)\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)} = 0.$$

Moreover  $u$  is stable : there is  $\varepsilon > 0$  such that if  $\|u|_{t=0} - v_0\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)} \leq \varepsilon$  then there is a **unique global solution** associated with  $v_0$ .

**Remarks.**

- The same result holds in the more general framework of  $BMO^{-1}$  [Auscher, Dubois, Tchamitchian '04].
- The result shows that  $\mathcal{G}$  is **open** in the strong topology. An immediate corollary of the theorem is that  $\mathcal{G}$  is **connected**.

# Idea of the proof of the result (large time behaviour)

**An easy case :** assume  $u_0 := u|_{t=0} \in L^2 \cap \dot{H}^{\frac{1}{2}}(\mathbb{R}^3)$ . Then  $u$  satisfies the energy inequality and in particular  $u \in L^4(\mathbb{R}^+; \dot{H}^{\frac{1}{2}}(\mathbb{R}^3))$  so there is  $t_0$  such that  $\|u(t_0)\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)} \leq \varepsilon_0$  and then that holds for all  $t \geq t_0$  by small data theory.

**The general case :** write  $u_0 = v_0 + w_0$  with  $w_0$  small in  $\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)$  and  $v_0$  in  $L^2(\mathbb{R}^3)$ .

Solve (NS) globally with the data  $w_0$ , the solution  $w(t)$  remains small in  $\dot{H}^{\frac{1}{2}}(\mathbb{R}^3)$  for all times.

Prove that the solution of

$$\partial_t v + \mathbb{P}(-v \cdot \nabla v + v \cdot \nabla w + w \cdot \nabla v) - \Delta v = 0$$

is bounded in the energy space and conclude as above.

# The set $\mathcal{G}$ is weakly open

We consider sequences **converging in the sense of distributions** to an element of  $\mathcal{G}$ .

**Examples** : the sequence  $\phi_n(x) := 2^n \phi(2^n x)$  converges weakly to zero. If  $\mathcal{G}$  were open for the weak topology then  $\phi$  would belong to  $\mathcal{G}$  by scale invariance... The same goes for  $\tilde{\phi}_n(x) := \phi(x - x_n)$ ,  $|x_n| \rightarrow \infty$ .

Define  $\Delta_k^h$  and  $\Delta_j^v$  Littlewood-Paley frequency truncation operators :

$$\mathcal{F}(\Delta_k^h f)(\xi) := \varphi(2^{-k} |(\xi_1, \xi_2)|) \mathcal{F}(f)(\xi)$$

$$\mathcal{F}(\Delta_j^v f)(\xi) := \varphi(2^{-j} |\xi_3|) \mathcal{F}(f)(\xi)$$

where  $\varphi \in \mathcal{C}_c^\infty(\frac{1}{2}, 1)$ , so that  $\sum_k \Delta_k^h f = \sum_j \Delta_j^v f = f$ . Notice that

$$\|\Delta_k^h \partial_1 f\|_{L^p} \sim 2^k \|\Delta_k^h f\|_{L^p}.$$

Then consider the norm  $\|f\|_{\mathcal{B}_q^1} := \left( \sum_{j,k} 2^{(j+k)q} \|\Delta_k^h \Delta_j^v f\|_{L^1(\mathbb{R}^3)}^q \right)^{\frac{1}{q}}$ .

**Remark** : scale invariance of (NS).



# The set $\mathcal{G}$ is weakly open

## Definition

Let  $0 < q \leq \infty$  be given. We say that a sequence  $(f_n)_{n \in \mathbb{N}}$ , bounded in  $\mathcal{B}_q^1$ , is **anisotropically oscillating** if the following property holds : for all sequences  $(k_n, j_n) \in \mathbb{Z}^{\mathbb{N}} \times \mathbb{Z}^{\mathbb{N}}$ ,

$$\limsup_{n \rightarrow \infty} 2^{j_n + k_n} \|\Delta_{k_n}^h \Delta_{j_n}^v f_n\|_{L^1} = C > 0 \implies \lim_{n \rightarrow \infty} |j_n - k_n| = \infty.$$

**Example** : the sequence

$$\phi_n(x) := 2^{\alpha n} \phi(2^{\alpha n} x_1, 2^{\alpha n} x_2, 2^{\beta n} x_3), \quad \alpha \neq \beta$$

is anisotropically oscillating : horizontal frequencies  $\sim 2^{\alpha n}$  and vertical frequencies  $\sim 2^{\beta n}$  so

$$\limsup_{n \rightarrow \infty} 2^{j_n + k_n} \|\Delta_{k_n}^h \Delta_{j_n}^v \phi_n\|_{L^1} = C > 0 \implies k_n \sim \alpha n, j_n \sim \beta n.$$

# The set $\mathcal{G}$ is weakly open

**Theorem** [Bahouri-G '12, Bahouri-Chemin-G in progress]

Let  $q \in ]0, 1[$  be given and let  $(u_{0,n})_{n \in \mathbb{N}}$  be a sequence of divergence free vector fields bounded in  $\mathcal{B}_q^1$ , converging towards  $u_0 \in \mathcal{B}_q^1$  in the sense of distributions, with  $u_0 \in \mathcal{G}$ . If  $u_0 - (u_{0,n})_{n \in \mathbb{N}}$  is anisotropically oscillating, then up to extracting a subsequence,  $u_{0,n} \in \mathcal{G}$  for all  $n \in \mathbb{N}$ .

## Remarks.

- One can essentially consider **any** bounded sequence **except** for sequences of the type described above and their superpositions.
- The theorem may be generalized by adding two more sequences to  $(u_{0,n})_{n \in \mathbb{N}}$ , where in each additional sequence the “privileged” direction is not  $x_3$  but  $x_1$  or  $x_2$ .
- The same result holds for data not in  $\mathcal{G}$ , on some life span  $[0, T]$  for  $T < T^*$ .

The rest of the talk is devoted to a sketch of the proof of this result.

- 1 Write down an “**anisotropic profile decomposition**” of the sequence of initial data. This allows to replace the sequence, up to an arbitrarily small remainder term, by a finite (large) sum of profiles of the type

$$\frac{1}{\lambda_n} \Phi\left(\frac{x_1}{\lambda_n}, \frac{x_2}{\lambda_n}, \frac{h_n x_3}{\lambda_n}\right) \quad h_n \rightarrow 0.$$

- 2 Propagate **globally in time** by (NS) each individual profile of the decomposition.
- 3 Prove that the construction of the previous step does provide, after **superposition** of all the global solutions, an **approximate solution** to the Navier-Stokes equations.

Before carrying out that program we shall **discuss an example** of the type above.

# A (typical) example (I)

Consider the divergence-free initial data

$$\Phi_{0,n}(x) := \frac{1}{\lambda_n} (\Phi_0^1, \Phi_0^2, 0) \left( \frac{x_1}{\lambda_n}, \frac{x_2}{\lambda_n}, \frac{h_n x_3}{\lambda_n} \right), \quad h_n \rightarrow 0.$$

Up to rescaling by  $\lambda_n$  it is equivalent to study

$$\tilde{\Phi}_{0,n}^h(x) := \Phi_0^h(x_h, h_n x_3), \quad x_h := (x_1, x_2), \quad \Phi_0^h := (\Phi_0^1, \Phi_0^2).$$

## A (typical) example (II)

Recall

$$\tilde{\Phi}_{0,n}^h(x) := \Phi_0^h(x_h, h_n x_3), \quad x_h := (x_1, x_2), \quad \Phi_0^h := (\Phi_0^1, \Phi_0^2).$$

To prove there is a **unique global solution** to (NS) associated with  $\tilde{\Phi}_{0,n}^h$  for  $n$  large enough [Chemin-G '10], we start by solving globally the **two dimensional equations** with data  $\Phi_0^h(x_h, y_3)$  for each  $y_3$ . We denote the solution by  $\Phi^h(t, x_h, y_3)$ .

Then we check that  $(\Phi^h, 0)(t, x_h, h_n x_3)$  is a global **approximate solution** to (NS) with data  $\tilde{\Phi}_{0,n}$ , so by rescaling, a global approximate solution associated with  $\Phi_{0,n}$  is

$$\Phi_n(t, x) := \frac{1}{\lambda_n} (\Phi^h, 0) \left( \frac{t}{\lambda_n^2}, \frac{x_h}{\lambda_n}, \frac{h_n x_3}{\lambda_n} \right).$$

## Another (typical) example

In the previous example

$$\Phi_{0,n}(x) := \frac{1}{\lambda_n}(\Phi_0^1, \Phi_0^2, 0)\left(\frac{x_1}{\lambda_n}, \frac{x_2}{\lambda_n}, \frac{h_n x_3}{\lambda_n}\right), \quad h_n \rightarrow 0,$$

we had  $\Phi_{0,n}^h \rightharpoonup 0$  if  $\lambda_n \rightarrow 0$  or  $\infty$  and  $\Phi_{0,n}^h(x) \rightharpoonup \Phi_0^h(x_h, 0)$  if  $\lambda_n \equiv 1$ .

Consider now the divergence-free initial data

$$u_{0,n} := u_0 + (\Phi_0^h, 0)(x_1, x_2, h_n x_3),$$

with  $u_0 \in \mathcal{G}$ . We assume that  $u_{0,n} \rightharpoonup u_0$  so  $(\Phi_0^h, 0)(x_h, 0) \equiv 0$ .

We know there is a global solution to (NS) associated with  $(\Phi_0^h, 0)(x_1, x_2, h_n x_3)$ , denoted  $\Phi_n(t, x)$ , and we call  $u$  the global solution associated with  $u_0$ . We want to prove that  $u + \Phi_n$  is a global, approximate solution to (NS).

## Another (typical) example

Since  $(\Phi_0^h, 0)(x_h, 0) \equiv 0$ , then  $\Phi_n(t, x_h, 0) \sim 0$  so up to a small error, the support in  $x_3$  of  $\Phi_n$  is  $\sim h_n^{-1} \rightarrow \infty$ .

Approximating  $u$  by a compactly supported vector field we find that the supports of  $u$  and  $\Phi_n$  are asymptotically disjoint, so the two vector fields **do not interact**.

That ends the proof in this model case.

# Profile decompositions

- Introduced by Gérard ('96), to describe the **lack of compactness** in  $\dot{H}^s(\mathbb{R}^d) \hookrightarrow L^p(\mathbb{R}^d)$ , extended to other Sobolev and Besov spaces by Jaffard ('99), Koch ('11), and to more general situations by Bahouri, Cohen and Koch ('11).

See also Brézis-Coron ('85), Métivier-Schochet ('98), Tintarev et al. ('07).

- **Applications to nonlinear PDEs** : among others by Merle and Vega ('98), Bahouri and Gérard ('99), Keraani ('01), G ('01), Bégout and Vargas ('07), Kenig and Merle ('08), Kenig and Koch ('10), G, Koch and Planchon ('12)...



# Anisotropic profile decomposition

Define  $\Lambda_{\lambda_n} \phi(x) := \frac{1}{\lambda_n} \phi\left(\frac{x}{\lambda_n}\right)$ ,  $\tilde{\Lambda}_{\lambda_n} \tilde{\phi}(t, x) := \frac{1}{\lambda_n} \tilde{\phi}\left(\frac{t}{\lambda_n^2}, \frac{x}{\lambda_n}\right)$ .

We say that  $\lambda_n^1 \perp \lambda_n^2$  if  $\lim_{n \rightarrow \infty} \left( \frac{\lambda_n^1}{\lambda_n^2} + \frac{\lambda_n^2}{\lambda_n^1} \right) = \infty$ .

In the following we use the notation  $[f]_\varepsilon(x) := f(x_1, x_2, \varepsilon x_3)$ .

**Theorem** [Bahouri-G '12, Chemin-Bahouri-G]

Let  $(u_n)_{n>0}$  be bounded in  $\mathcal{B}_q^1$ ,  $q < 1$ . Then, up to an extraction,

$$\begin{aligned} u_{0,n} &= u_0 + \left[ (v_n^{0,h} + h_n^0 w_n^{0,h}, w_n^{0,3}) \right]_{h_n^0} \\ &\quad + \sum_{j=1}^L \Lambda_{\lambda_n^j} \left[ (v_n^{j,h} + h_n^j w_n^{j,h}, w_n^{j,3}) \right]_{h_n^j} + \rho_n^L \end{aligned}$$

with  $(h_n^j)_{n \in \mathbb{N}} \rightarrow 0$  as  $n \rightarrow \infty$ , and  $(\lambda_n^j)_{j \geq 1}$  are mutually orthogonal, going to zero or infinity as  $n \rightarrow \infty$ .

# Anisotropic profile decomposition

Moreover  $(v_n^{0,h}, 0)$  and  $w_n^0$  are smooth, divergence free vector fields, and satisfy the following bounds :

$$\sum_{j \in \mathbb{N}} (\|v_n^{j,h}\|_{\mathcal{B}_1^1} + \|w_n^{j,3}\|_{\mathcal{B}_1^1}) \leq C,$$
$$\|v_n^{0,h}(\cdot, 0)\|_{\dot{B}_{1,1}^1(\mathbb{R}^2)} + \|w_n^{0,3}(\cdot, 0)\|_{\dot{B}_{1,1}^1(\mathbb{R}^2)} \rightarrow 0.$$

Finally  $\rho_n^L$  is a “remainder”, in a sense that for some scale-invariant space  $\mathcal{X}$  containing  $\mathcal{B}_1^1$ ,

$$\limsup_{n \rightarrow \infty} \|\rho_n^L\|_{\mathcal{X}} \rightarrow 0, \quad L \rightarrow \infty.$$

# Time evolution of the decomposition

- Define  $\Phi_n^{0,0} := u_0 + [(v_n^{0,h} + h_n^0 w_n^{0,h}, w_n^{0,3})]_{h_n^0}$  and for  $j \geq 1$ ,  
 $\Phi_n^{j,0} := \Lambda_{\lambda_n^j} [(v_n^{j,h} + h_n^j w_n^{j,h}, w_n^{j,3})]_{h_n^j}$ . Then as in the model example, for  $n$  large enough,  $\Phi_n^{j,0}$  generates a unique, global solution  $\Phi_n^j(t)$  which can be written for  $j \geq 1$

$$\Phi_n^j = \tilde{\Lambda}_{\lambda_n^j} \tilde{\Phi}_n^j$$

with  $\tilde{\Phi}_n^j$  satisfying "good" bounds.

- The remainder term  $\rho_n$  can easily be evolved by the Navier-Stokes flow since it is arbitrarily small in scale-invariant spaces. We call  $\mathcal{R}_n$  the associate solution.

# Approximate solution

Define

$$w_n := u_n - \left( \sum_{0 \leq j \leq L} \Phi_n^j + \mathcal{R}_n \right),$$

and prove that  $w_n$  exists globally.

Indeed  $w_n$  solves a **perturbed (NS) equation** of the type

$$\partial_t w_n + w_n \cdot \nabla w_n + G_n \cdot \nabla w_n + w_n \cdot \nabla G_n - \Delta w_n = -H_n - \nabla p_n$$

with  $\operatorname{div} w_n = 0$  and initial data  $w_n|_{t=0} = 0$ .

So we need to prove that  $G_n$  satisfies uniform bounds in some adequate space  $\mathcal{Y}$  and that  $H_n$  is small in some other adequate space  $\mathcal{Y}'$ .

# The forcing term

It is enough to prove that for all  $j \neq k$ ,

$$\limsup_{n \rightarrow \infty} \|\tilde{\Lambda}_{\lambda_n^j}^n \tilde{\Phi}_n^j \otimes \tilde{\Lambda}_{\lambda_n^k}^n \tilde{\Phi}_n^k\|_{L^1(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} = 0.$$

Assume that  $\lambda_n^j / \lambda_n^k$  goes to zero as  $n$  goes to infinity. Then

$$\begin{aligned} \|\tilde{\Lambda}_{\lambda_n^j}^n \tilde{\Phi}_n^j \otimes \tilde{\Lambda}_{\lambda_n^k}^n \tilde{\Phi}_n^k\|_{L^1(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} &\leq \|\tilde{\Lambda}_{\lambda_n^j}^n \tilde{\Phi}_n^j\|_{L^1(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} \\ &\quad \times \|\tilde{\Lambda}_{\lambda_n^k}^n \tilde{\Phi}_n^k\|_{L^\infty(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} \end{aligned}$$

and we use the fact that

$$\|\tilde{\Lambda}_{\lambda_n^j}^n \tilde{\Phi}_n^j\|_{L^1(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} \lesssim \lambda_n^j \quad \text{and} \quad \|\tilde{\Lambda}_{\lambda_n^k}^n \tilde{\Phi}_n^k\|_{L^\infty(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} \lesssim \frac{1}{\lambda_n^k}.$$

So

$$\|\tilde{\Lambda}_{\lambda_n^j}^n \tilde{\Phi}_n^j \otimes \tilde{\Lambda}_{\lambda_n^k}^n \tilde{\Phi}_n^k\|_{L^1(\mathbb{R}^+; \dot{B}_{1,1}^{2,1})} \leq \frac{\lambda_n^j}{\lambda_n^k}$$

which proves the result.

# Extensions

- Add **geometry** (here spectral localization is about a plane).
- Use more deeply the **structure of divergence free vector fields**.
- What about the blow-up behaviour (if such solutions exist)? cf. for instance [Sverak-Rusin '12].