

# MAT Syllabus Practice Solutions

James Munro\*

updated July 2020 (first version October 2019)

## Polynomials

- Solve  $x^2 - x - 1 = 0$   
The quadratic formula gives  $x = \frac{1 \pm \sqrt{5}}{2}$ .
- Solve  $x^4 - x^2 - 1 = 0$   
Write  $y = x^2$  to get a quadratic for  $y$ . This is the quadratic above for  $y$ , so  $x^2 = \frac{1 \pm \sqrt{5}}{2}$ . But  $x^2 \geq 0$  so  $x = \pm \sqrt{\frac{1 \pm \sqrt{5}}{2}}$ .
- Write  $x^2 + 4x + 3$  in the form  $(x + a)^2 + b$   
 $(x + 2)^2 - 1$
- How many real solutions does  $x^2 + bx + 1 = 0$  have? Find the different cases in terms of  $b$ .  
The discriminant,  $b^2 - 4$ , is positive if  $b > 2$  or  $b < -2$ , negative if  $-2 < b < 2$  and zero if  $b = \pm 2$ . So there are two real solutions if  $b > 2$  or if  $b < -2$ , one real solution if  $b = \pm 2$  and no real solutions otherwise.
- Factorise  $x^2 + 4x + 3$   
 $(x + 1)(x + 3)$
- Let  $p(x) = x^3 - 13x^2 - 65x - 51$ . Check that  $p(17) = 0$ . Factorise  $p(x)$ .  
 $p(17) = 17^3 - 13 \times 17^2 - 65 \times 17 - 51 = 17(17^2 - 13 \times 17 - 65 - 3) = 17^2(17 - 13 - 4) = 0$ .  
So  $(x - 17)$  is a factor. Polynomial division gives  $p(x) = (x - 17)(x^2 + 4x + 3)$ , so  $p(x) = (x - 17)(x + 1)(x + 3)$ .

## Algebra

- Solve the simultaneous equations  $x + y = 1$  and  $x - y = 3$ .  
 $x = 2$  and  $y = -1$ .
- For which values of  $x$  is it true that  $x^2 + 4x + 3 > 0$ ?  
 $x > -1$  or  $x < -3$ .
- Expand  $(2x + 3)^3$   
 $8x^3 + 36x^2 + 54x + 27$
- I've got four playing cards; the ace and king of clubs, and the ace and king of hearts. I shuffle the cards together and deal them out left to right. What's the probability that the kings and aces alternate? (they alternate if they are either arranged as  $AKAK$  or  $KAKA$ )  
There are 24 possible orders for the cards. Eight of these have alternating kings and aces, so the probability is  $1/3$ .

## Differentiation

- Differentiate  $x^{17}$  with respect to  $x$ .  
 $17x^{16}$

---

\*Admissions Coordinator for Maths at Oxford. Comments or corrections should be sent to [james.munro@maths.ox.ac.uk](mailto:james.munro@maths.ox.ac.uk)

- Differentiate  $\sqrt{x}$  with respect to  $x$ .

$$\frac{1}{2\sqrt{x}}$$

- Differentiate  $e^{3x}$  with respect to  $x$ .

$$3e^{3x}$$

- Differentiate  $2e^{-x} - x^2$  with respect to  $x$ .

$$-2e^{-x} - 2x$$

- Find the tangent to the curve  $y = e^x + 1$  at  $x = 2$ .

$$y = e^2(x - 2) + e^2 + 1$$

- Find the normal to the parabola  $y = x^2$  at  $x = 3$ .

$$y = -\frac{1}{6}(x - 3) + 9$$

- Find the turning points of the curve  $y = x^4 - 2x^3 + x^2$ . Identify whether the turning points are maxima or minima.

Turning points at  $x = 0$  (minimum),  $x = \frac{1}{2}$  (maximum),  $x = 1$  (minimum).

- For which values of  $x$  is  $y = x^4 - 2x^3 + x^2$  increasing? For which values of  $x$  is it decreasing?

Increasing for  $0 < x < \frac{1}{2}$  and for  $1 < x$ . Decreasing for  $x < 0$  and for  $\frac{1}{2} < x < 1$ .

- Two points  $A$  and  $B$  are on the curve  $y = x^3 + x^2 + x + 1$ .  $A$  is held fixed at  $(1, 4)$ . The point  $B$  is moved along the curve towards  $A$ . What happens to the line through  $A$  and  $B$ ?

The tangent at  $A$  is  $y = 6x - 2$ . If the line  $AB$  has equation  $y = mx + c$  say, then  $m$  gets closer and closer to 6 and  $c$  gets closer and closer to  $-2$ .

### Integration

- Suppose that the derivative of a polynomial  $p(x)$  with respect to  $x$  is  $q(x)$ . Find  $\int q(x) dx$ .

$p(x) + c$  where  $c$  is a constant

- Find the area enclosed by the polynomial  $x^2 + 4x + 3 = 0$  and the  $x$ -axis.

$$\frac{4}{3}$$





- Find  $\int_{-1}^1 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 dx$

Note that  $\int_{-1}^1 x^a dx = 0$  for  $a$  odd. The integral is  $2(1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7}) = \frac{352}{105}$ .


### Graphs

- Sketch graphs of

$$y = x^2 + 4x + 3, \quad y = x^3 + 4x^2 + 3x, \quad y = 2^x, \quad y = \log_2 x \quad \text{on separate axes.}$$



- Sketch graphs of  $y = \sin x$ ,  $y = \cos x$  and  $y = \tan x$  on the same axes.



### Logarithms and powers

- Simplify  $\log 3 + \log 4$  into a single term.

$$\log 12$$

- Expand  $(e^x + e^{-x})(e^x + e^{-x})$


$$e^{2x} + 2 + e^{-2x}.$$

- Solve  $2^x = 3$ .

$$x = \log_2 3$$

### Transformations

- Let  $f(x) = x^2 + 4x + 3$ . If you didn't sketch a graph of this before, sketch one now.
- Sketch a graph of  $y = f(x + 2)$ .
- Sketch a graph of  $y = 3f(2x)$ .



### Geometry

- Add the vectors  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  and  $\begin{pmatrix} 3 \\ -2 \end{pmatrix}$ .

$$\begin{pmatrix} 4 \\ 0 \end{pmatrix}$$

- Find the equation of the line through  $(1, 0)$  and  $(0, -1)$ .

$$y = x - 1$$

- Find the equation of the line through  $(1, 2)$  with gradient 3.

$$y = 3(x - 1) + 2 = 3x - 1$$

- A circle has centre  $(-1, 4)$  and radius 3. Write down an equation for the circle.

$$(x + 1)^2 + (y - 4)^2 = 9$$

- What's the area of this circle?

$$9\pi$$

- Points  $A$  and  $B$  lie on a circle with centre  $O$  and radius 1. The angle  $\angle AOB$  is  $120^\circ$ . Find the length of the arc between  $A$  and  $B$ . Find the area enclosed by that arc and the radii  $OA$  and  $OB$ .

It's a third of a circle, so the arc length is  $2\pi/3$  and the area is  $\pi/3$ .

## Trigonometry

- Solve  $\sin x = \frac{1}{2}$ .  
 $x = 30^\circ + n \times 360^\circ$ , or  $x = 150^\circ + n \times 360^\circ$ , for any whole number  $n$ .
- Solve  $\tan x = 1$ .  
 $x = 45^\circ + n \times 180^\circ$  for any whole number  $n$
- Write  $\cos^4 x + \cos^2 x$  in terms of  $\sin x$ .  
 $(1 - \sin^2 x)^2 + (1 - \sin^2 x) = 2 - 3\sin^2 x + \sin^4 x$ .
- Simplify  $\cos(450^\circ - x)$   
 $\sin x$
- A triangle  $ABC$  has side lengths  $AB = 3$  and  $BC = 2$ , and the angle  $\angle ABC = 120^\circ$ . Find the remaining side length  $AC$ , the area of the triangle, and an expression for  $\sin \angle BCA$ .  
Cosine rule;  $AC = \sqrt{19}$ . The area of the triangle is  $3\sqrt{3}/2$ . Sine rule;  $\sin \angle BCA = (3\sqrt{3}/2\sqrt{19})$

## Sequences and series

- A sequence is defined by  $a_0 = 1$ ,  $a_1 = 1$ ,  $a_2 = 1$ , and

$$a_n = a_{n-1} + a_{n-2} + a_{n-3} \quad \text{for } n \geq 3.$$

Find  $a_{10}$ .

$$a_3 = 3, a_4 = 5, a_5 = 9, a_6 = 17, a_7 = 31, a_8 = 57, a_9 = 105, a_{10} = 193.$$

- A sequence has first term 3 and each subsequent term is 5 more than the previous term. Find the sum of the first four terms.  
 $4 \times 3 + \frac{4 \times 3}{2} \times 5 = 42$
- A sequence has first term 4 and each subsequent term is 6 times more than the previous term. Find the sum of the first four terms.  
 $4(1 + 6 + 6^2 + 6^3) = 4 \frac{6^4 - 1}{6 - 1} = 4 \frac{1295}{5} = 4 \times 259 = 1036$ .
- When does the sum  $1 + x^3 + x^6 + x^9 + x^{12} + \dots$  converge? Simplify it in the case that it converges.  
Converges when  $-1 < x < 1$ . In that case, it converges to  $1/(1 - x^3)$ .